

(2 + 1)-dimensional topological quantum field theory from subfactors and Dehn surgery formula for 3-manifold invariants

YASUYUKI KAWAHIGASHI*

Department of Mathematical Sciences
University of Tokyo, Komaba, Tokyo, 153-8914, JAPAN
e-mail: yasuyuki@ms.u-tokyo.ac.jp

NOBUYA SATO*

Department of Mathematics and Information Sciences
Osaka Prefecture University, Sakai, Osaka, 599-8531, JAPAN
e-mail: nobuya@mi.cias.osakafu-u.ac.jp

MICHIHISA WAKUI

Department of Mathematics
Osaka University, Toyonaka, Osaka, 560-0043, JAPAN
e-mail: wakui@math.sci.osaka-u.ac.jp

February 1, 2008

Abstract

In this paper, we establish the general theory of (2+1)-dimensional topological quantum field theory (in short, TQFT) with a Verlinde basis. It is a consequence that we have a Dehn surgery formula for 3-manifold invariants for this kind of TQFT's. We will show that Turaev-Viro-Ocneanu unitary TQFT's obtained from subfactors satisfy the axioms of TQFT's with Verlinde bases. Hence, in a Turaev-Viro-Ocneanu TQFT, we have a Dehn surgery formula for 3-manifolds. It turns out that this Dehn surgery formula is nothing but the formula of the Reshetikhin-Turaev invariant constructed from a tube system, which is a modular category corresponding to the quantum double construction of a C^* -tensor category. In the forthcoming paper, we

*Supported in part by the Grants-in-Aid for Scientific Research, JSPS.

will exhibit computations of Turaev-Viro-Ocneanu invariants for several “basic 3-manifolds”. In Appendix, we discuss the relationship between the system of M_∞ - M_∞ bimodules arising from the asymptotic inclusion $M \vee M^{op} \subset M_\infty$ constructed from $N \subset M$ and the tube system obtained from a subfactor $N \subset M$.

1 Introduction

Since the discovery of the celebrated Jones polynomial [9], there have been a great deal of studies on quantum invariants of knots, links, and 3-manifolds. Among them, we deal with the two constructions in this paper; Ocneanu’s generalization of the Turaev-Viro invariants [22] of 3-manifolds based on triangulation and the Reshetikhin-Turaev invariants [16] of those based on Dehn surgery. The former uses quantum $6j$ -symbols arising from subfactors and the tensor categories we have are not necessarily braided, while the latter invariants require braiding, or more precisely, modular tensor categories. Methods to construct a modular tensor category from a rational tensor category have been studied by several authors and are often called “quantum double” constructions. In the setting of subfactor theory, such a method was first found by Ocneanu using his generalization of the Turaev-Viro topological quantum field theory (TQFT) and Izumi [7] later gave a formulation based on sector theory.

We start with a (rational unitary) tensor category (arising from a subfactor), which is not braided in general. Then we obtain two TQFT’s with the methods above as follows. One is a Turaev-Viro-Ocneanu TQFT based on a state sum and triangulation which directly uses $6j$ -symbols of the tensor category. The other is a Reshetikhin-Turaev TQFT arising from the modular tensor category we obtain with the quantum double construction of the original category. It is natural to ask what relation we have between these two TQFT’s. In order to make a general study on such TQFT’s, we first establish a Dehn surgery formula (Proposition 2.3) for a general TQFT, which represents an invariant of a 3-manifold in terms of a weight sum of invariants of links that we use for the Dehn surgery construction of the 3-manifold. This formula uses a basis of the Hilbert space for $S^1 \times S^1$ arising from the TQFT. We show that several nice properties for such a basis are mutually equivalent (Theorem 2.10) and we say that a basis is a *Verlinde basis* when such properties hold (Definition 2.5). We next show that a Turaev-Viro-Ocneanu TQFT has a Verlinde basis in this sense (Theorem 4.6) and then as a corollary of the Dehn surgery formula, we conclude that the above two TQFT’s are identical for any closed 3-manifold (Theorem 5.2). (Actually, the claim that a Turaev-Viro-Ocneanu TQFT has a Verlinde basis has been announced by Ocneanu and a proof is presented in [2, Chapter 12], but normalizations are inaccurate there, so we include a proof for this claim for the sake of completeness here, along the line of Ocneanu’s tube algebra and braiding arising from the Turaev-Viro-Ocneanu TQFT.) This identity result has been announced by

two of us in [18]. It has been also announced in page 244 of Ocneanu [15], but it seems to us that his line of arguments are different from ours which relies on our Dehn surgery formula. This result also proves a conjecture in [13, Section 8.2] and gives an answer to a question in [10, page 546]. We refer to the book [2] for subfactor theory and its applications. Throughout this paper, we only consider subfactors with finite depth and finite index.

Acknowledgement: A part of this work was done while the first two authors visited Mathematical Sciences Research Institute at Berkeley. We appreciate their financial supports and hospitality.

2 $(2+1)$ -dimensional TQFT with Verlinde basis and Dehn surgery formula

Giving a $(2+1)$ -dimensional topological quantum field theory Z , we have a 3-manifold invariant $Z(M)$ in the canonical way. The purpose of this section is to describe $Z(M)$ in terms of the S -matrix and framed links, and to give a criterion for the Verlinde identity [23] to hold in our framework.

Throughout this section, we use the following notations. For an oriented manifold M , we denote by $-M$ the same manifold with the opposite orientation. For an orientation preserving diffeomorphism $f : M \rightarrow N$, we denote by $-f$ the same diffeomorphism viewed as an orientation preserving diffeomorphism $-M \rightarrow -N$. We regard the empty set $\emptyset$ as an oriented closed surface with the unique orientation. The dual vector space of a finite dimensional vector space V over $\mathbb{C}$ is denoted by V^* , and the dual basis of a basis $\{v_i\}_i$ is denoted by $\{v_i^*\}_i$. The closure of a subspace X in the 3-sphere S^3 is denoted by $\overline{X}$.

Convention of orientations for manifolds.

Throughout this section, we assume that the 3-sphere S^3 is oriented, and assume that any 3-dimensional submanifold of S^3 is oriented by the orientation induced from S^3 . We also assume that the solid torus $D^2 \times S^1$ equips with the orientation such that the diffeomorphism $h : D^2 \times S^1 \rightarrow \mathbb{R}^3 \subset S^3$ defined by

$$h((x, y), e^{i\theta}) = ((2+x)\cos\theta, (2+x)\sin\theta, y)$$

is orientation preserving, and assume that the torus $S^1 \times S^1$ is oriented by the orientation induced from $D^2 \times S^1$ (see Figure 1).

Let us recall the definition of $(2+1)$ -dimensional topological quantum field theory due to Atiyah [1], [5]. By a 3-cobordism we mean a triple $(M; \Sigma_1, \Sigma_2)$ consisting of a compact oriented 3-manifold M and two closed oriented surfaces Σ_1, Σ_2 such that $\partial M = (-\Sigma_1) \cup \Sigma_2$ and $\Sigma_1 \cap \Sigma_2 = \emptyset$. For an oriented closed surface Σ , the identity cobordism is given by $Id_\Sigma = (\Sigma \times [0, 1]; \Sigma \times \{0\}, \Sigma \times \{1\})$.




Figure 1: the orientation of torus $S^1 \times S^1$

Definition 2.1. *Let Z be a functor consisting of the following three functions.*

- (1) *To each closed oriented surface Σ , it assigns a finite dimensional $\mathbb{C}$ -vector space $Z(\Sigma)$.*
- (2) *To each cobordism W , it assigns a $\mathbb{C}$ -linear map Z_W .*
- (3) *To each orientation preserving diffeomorphism f between closed oriented surfaces, it assigns a $\mathbb{C}$ -linear isomorphism $Z(f)$.*

If Z has the following properties, it is called a $(2+1)$ -dimensional topological quantum field theory (TQFT in short).

- (i) *Z is functorial with respect to the composition of orientation preserving diffeomorphisms. More precisely,*
 - (a) *$Z(g \circ f) = Z(g) \circ Z(f)$ for orientation preserving diffeomorphisms f and g .*
 - (b) *$Z(id_\Sigma) = id_{Z(\Sigma)}$ for an oriented closed surface Σ .*
- (ii) *Z is functorial with respect to the composition of 3-cobordisms. More precisely,*
 - (a) *If two cobordisms $W_1 = (M_1; \Sigma_1, \Sigma_2)$ and $W_2 = (M_2; \Sigma_2, \Sigma_3)$ are obtained by cutting a cobordism $W = (M; \Sigma_1, \Sigma_3)$ along Σ_2 in $M = M_1 \cup M_2$, then*

$$Z_W = Z_{W_2} \circ Z_{W_1}.$$

- (b) *$Z(i_1)^{-1} \circ Z_{Id_\Sigma} \circ Z(i_0) = id_{Z(\Sigma)}$ for the identity cobordism on Σ , where $i_t : \Sigma \longrightarrow \Sigma \times \{t\}$ ($t = 0, 1$) are orientation preserving diffeomorphisms defined by $i_t(x) = (x, t)$, $x \in \Sigma$.*
- (iii) *Let $W = (M, \Sigma_1, \Sigma_2)$, $W' = (M', \Sigma'_1, \Sigma'_2)$ be two cobordisms. Suppose that there is an orientation preserving diffeomorphism $h : M \longrightarrow M'$*

such that $h(\Sigma_i) = \Sigma'_i$ for $i = 1, 2$. Then, for $f_1 := -h|_{\Sigma_1} : \Sigma_1 \longrightarrow \Sigma'_1$ and $f_2 := h|_{\Sigma_2} : \Sigma_2 \longrightarrow \Sigma'_2$, the following diagram commutes.

$$\begin{array}{ccc} Z(\Sigma_1) & \xrightarrow{Z(f_1)} & Z(\Sigma'_1) \\ Z_W \downarrow & & \downarrow Z_{W'} \\ Z(\Sigma_2) & \xrightarrow{Z(f_2)} & Z(\Sigma'_2) \end{array}$$

(iv) Let $W_1 = (M; \Sigma_1, \Sigma_2)$, $W_2 = (N; \Sigma'_2, \Sigma_3)$ be two cobordisms, and $f : \Sigma_2 \longrightarrow \Sigma'_2$ an orientation preserving diffeomorphism. Then, for the cobordism $W := (N \cup_f M; \Sigma_1, \Sigma_3)$ obtained by gluing of W_1 to W_2 along f ,

$$Z_W = Z_{W_2} \circ Z(f) \circ Z_{W_1}.$$

(v) There are natural isomorphisms

- (a) $Z(\emptyset) \cong \mathbb{C}$.
- (b) $Z(\Sigma_1 \amalg \Sigma_2) \cong Z(\Sigma_1) \otimes Z(\Sigma_2)$ for oriented closed surfaces Σ_1 and Σ_2 .
- (c) $Z(-\Sigma) \cong Z(\Sigma)^*$ for an oriented closed surface Σ .

Remarks 2.2. 1. For an oriented closed 3-manifold M , we have a cobordism $W = (M; \emptyset, \emptyset)$. This cobordism W induces a linear map

$$\mathbb{C} \cong Z(\emptyset) \xrightarrow{Z_W} Z(\emptyset) \cong \mathbb{C}.$$

We denote by $Z(M)$ the image of 1 under the above map. By the condition (iii) we see that $Z(M)$ is a topological invariant of M .

2. Let Γ_Σ denote the mapping class group of the oriented closed surface Σ . Then, we have a representation of Γ_Σ

$$\rho : \Gamma_\Sigma \longrightarrow GL(Z(\Sigma)), [f] \longmapsto Z(f),$$

where $[f]$ denotes the isotopy class of f .

3. For an oriented closed surface Σ , we have $\dim Z(\Sigma) = Z(\Sigma \times S^1)$.

4. By a 3-cobordism with parametrized boundary we mean a triple $(M; j_1, j_2)$ consisting of a compact oriented 3-manifold M , an orientation reversing embedding $j_1 : \Sigma_1 \longrightarrow \partial M$ and an orientation preserving embedding $j_2 : \Sigma_2 \longrightarrow \partial M$ such that $\partial M = (-j_1(\Sigma_1)) \cup j_2(\Sigma_2)$ and $j_1(\Sigma_1) \cap j_2(\Sigma_2) = \emptyset$. Any cobordism with parametrized boundary $\mathcal{W} = (M; \Sigma_1 \xrightarrow{j_1} \partial M, \Sigma_2 \xrightarrow{j_2} \partial M)$ induces a linear map $Z_{\mathcal{W}} : Z(\Sigma_1) \longrightarrow Z(\Sigma_2)$ such that the following diagram commutes.

$$\begin{array}{ccc}
Z(\Sigma_1) & \xrightarrow{Z_W} & Z(\Sigma_2) \\
Z(j_1) \downarrow & & \downarrow Z(j_2) \\
Z(j_1(\Sigma_1)) & \xrightarrow{Z_W} & Z(j_2(\Sigma_2))
\end{array}$$

Here, we set $W = (M; j_1(\Sigma_1), j_2(\Sigma_2))$.

Let Z be a $(2+1)$ -dimensional TQFT. We consider the cobordism $W := (Y \times S^1; \Sigma_1 \sqcup \Sigma_2, \Sigma_3)$, where Y is the compact oriented surface in $\mathbb{R}^3$ depicted in Figure 2 and $\Sigma_i = C_i \times S^1$ for $i = 1, 2, 3$. Then, W induces a linear map $Z_W : Z(S^1 \times S^1) \otimes Z(S^1 \times S^1) \longrightarrow Z(S^1 \times S^1)$. It can be easily verified that the map Z_W gives an associative algebra structure on $Z(S^1 \times S^1)$. The identity element of this algebra is given by $Z_{W_0}(1)$, where $W_0 := (D^2 \times S^1; \emptyset, S^1 \times S^1)$. We call this algebra the *fusion algebra associated with Z* .




Figure 2: the compact oriented surface Y

Let us introduce a Dehn surgery formula of $Z(M)$. Let Z be a $(2+1)$ -dimensional TQFT, and $S : S^1 \times S^1 \longrightarrow S^1 \times S^1$ the orientation preserving diffeomorphism defined by $S(z, w) = (\bar{w}, z)$, $(z, w) \in S^1 \times S^1$, where we regard S^1 as the set of complex numbers of absolute value 1. Given a basis $\{v_i\}_{i=0}^m$ of $Z(S^1 \times S^1)$, we define $S_{ji} \in \mathbb{C}$ by $Z(S)v_i = \sum_{j=0}^m S_{ji}v_j$.

Let $\{v_i\}_{i=0}^m$ be a basis of $Z(S^1 \times S^1)$. Then, we can define a framed link invariant as follows. Let $L = L_1 \cup \cdots \cup L_r$ be a framed link with r -components in the 3-sphere S^3 , and $h_i : D^2 \times S^1 \longrightarrow N(L_i)$ be the framing of L_i for each $i \in \{1, \dots, r\}$, where $N(L_i)$ denotes the tubular neighborhood of L_i . We fix an orientation for $\partial N(L_i)$ such that $j_i := h_i|_{\partial D^2 \times S^1} : S^1 \times S^1 \longrightarrow \partial N(L_i)$ is orientation preserving. Since the orientation for $N(L_i)$ is not compatible with the orientation for the link exterior $X := S^3 - N(L_1) \cup \cdots \cup N(L_r)$, we can consider the cobordism with parametrized boundary $\mathcal{W}_L := (X; \coprod_{i=1}^r j_i, \emptyset)$.

This cobordism induces a $\mathbb{C}$ -linear map $Z_{\mathcal{W}_L} : \bigotimes_{i=1}^r Z(S^1 \times S^1) \longrightarrow \mathbb{C}$. It is easy to see that for each $i_1, \dots, i_r = 0, 1, \dots, m$ the complex number $J(L; i_1, \dots, i_r) := Z_{\mathcal{W}_L}(v_{i_1} \otimes \cdots \otimes v_{i_r})$ is a framed link invariant of L .

Proposition 2.3. *Let Z be a $(2+1)$ -dimensional TQFT and $\{v_i\}_{i=0}^m$ a basis of $Z(S^1 \times S^1)$ such that v_0 is the identity element in the fusion algebra. Let M be a closed oriented 3-manifold obtained from S^3 by Dehn surgery along a framed link $L = L_1 \cup \cdots \cup L_r$. Then, the 3-manifold invariant $Z(M)$ is given by the formula*

$$Z(M) = \sum_{i_1, \dots, i_r=0}^m S_{i_1,0} \cdots S_{i_r,0} J(L; i_1, \dots, i_r).$$

Proof. Let X be the link exterior of L in S^3 , and $h_i : D^2 \times S^1 \longrightarrow \partial N(L_i)$ the framing of L_i for each $i \in \{1, \dots, r\}$. Then, by using attaching maps $f_i : S^1 \times S^1 \longrightarrow \partial N(L_i)$ satisfying $f_i(S^1 \times 1) = h_i(1 \times S^1)$, $i = 1, \dots, r$, we have

$$M = X \bigcup_{\coprod_{i=1}^r f_i} \left(\coprod_{i=1}^r D^2 \times S^1 \right).$$

Therefore,

$$Z(M) = Z_{W_2} \circ Z\left(\coprod_{i=1}^r f_i\right) \circ Z_{W_1},$$

where $W_1 := (\coprod^r (D^2 \times S^1); \emptyset, \coprod^r (S^1 \times S^1))$ and $W_2 := (X; -\partial X, \emptyset)$. This implies that

$$Z(M) = Z_{W_L} \circ \left(\bigotimes_{i=1}^r Z(j_i^{-1}) \right) \circ \left(\bigotimes_{i=1}^r Z(f_i) \right) \circ \left(\bigotimes_{i=1}^r Z_{W_0} \right),$$

where $j_i := h_i|_{\partial D^2 \times S^1} : S^1 \times S^1 \longrightarrow \partial N(L_i)$ for $i = 1, \dots, r$.

For each i we can choose f_i satisfying $j_i^{-1} \circ f_i = S$ up to isotopy as for f_i satisfying $f_i(S^1 \times 1) = h_i(1 \times S^1)$. Then, we have

$$Z(M) = Z_{W_L} \circ \left(\bigotimes_{i=1}^r Z(S) \right) \circ \left(\bigotimes_{i=1}^r Z_{W_0} \right).$$

This implies that

$$Z(M) = \sum_{i_1, \dots, i_r=0}^m S_{i_1,0} \cdots S_{i_r,0} J(L; i_1, \dots, i_r).$$

This completes the proof.

Q.E.D.

Proposition 2.4. *Let Z be a $(2+1)$ -dimensional TQFT and $\{v_i\}_{i=0}^m$ a basis of $Z(S^1 \times S^1)$. Let H be the Hopf link depicted as in Figure 3, and U the orientation preserving diffeomorphism from $S^1 \times S^1$ to $-S^1 \times S^1$ defined by $U(z, w) = (z, \bar{w})$. Then, the following are equivalent.*




Figure 3: the Hopf link H

(i) $J(H; i, j) = S_{ij}$ for all $i, j = 0, 1, \dots, m$.

(ii) $Z(U)v_i = v_i^*$ for all $i = 0, 1, \dots, m$.

Proof. Let $h_i : D^2 \times S^1 \longrightarrow N(H_i)$ be the framing of H_i for each $i = 1, 2$. Putting $j_i = h_i|_{S^1 \times S^1}$ ($i = 1, 2$) and $X := \overline{S^3 - N(H_1) \cup N(H_2)}$, we have a cobordism with parametrized boundary $\mathcal{W} := (X; j_1, -j_2)$. Then

$$Z_{\mathcal{W}}(v_i) = \sum_{j=0}^m J(H; i, j) v_j^*.$$

Let V_1 be the tubular neighborhood of $N(H_1)$ depicted as in Figure 4.




Figure 4:

Setting $V_2 := \overline{S^3 - V_1}$, and defining orientation preserving diffeomorphisms $j'_1 : S^1 \times S^1 \longrightarrow \partial V_1$ and $j'_2 : S^1 \times S^1 \longrightarrow \partial V_2$ in parallel to j_1 and j_2 respectively, we see that

- $\partial V_1 = -\partial V_2$,
- $X = \overline{V_1 - N(H_1) \cup V_2 - N(H_2)}$,
- $\mathcal{W}_1 := (\overline{V_1 - N(H_1)}; j_1, j'_1) \cong Id_{S^1 \times S^1}$ as cobordisms,
- $\mathcal{W}_2 := (\overline{V_2 - N(H_2)}; -j'_2, -j_2) \cong Id_{-S^1 \times S^1}$ as cobordisms.

Since

$$(-j'_2)^{-1} \circ j'_1 : \begin{cases} m \longmapsto -l, \\ l \longmapsto -m. \end{cases}$$




Figure 5:

for the meridian m and the longitude l (see Figure 5), it follows that

$$Z(-j_2'^{-1} \circ j_1') = Z(U \circ S).$$

Therefore, we obtain

$$Z_{\mathcal{W}} = Z_{\mathcal{W}_2} \circ Z(U \circ S) \circ Z_{\mathcal{W}_1},$$

and whence

$$\sum_{j=0}^m J(H; i, j) v_j^* = Z_{\mathcal{W}}(v_i) = (\theta \circ Z(U \circ S))(v_i),$$

where $\theta : Z(-S^1 \times S^1) \longrightarrow Z(S^1 \times S^1)^*$ denotes the natural isomorphism. Let $(U_{ij})_{i,j=0,1,\dots,m}$ be the presentation matrix of $\theta \circ Z(U) : Z(S^1 \times S^1) \longrightarrow Z(S^1 \times S^1)^*$ with respect to $\{v_i\}_{i=0}^m$ and $\{v_i^*\}_{i=0}^m$. Then, we have

$$(J(H; i, j))_{i,j=0,1,\dots,m} = (U_{ij})(S_{ij}).$$

Hence, we have

$$\begin{aligned} & (J(H; i, j))_{i,j=0,1,\dots,m} = (S_{ij})_{i,j=0,1,\dots,m} \\ \iff & (U_{ij}) = I, \text{ where } I \text{ is the identity matrix.} \\ \iff & (\theta \circ Z(U))(v_i) = v_i^* \ (i = 0, 1, \dots, m). \end{aligned}$$

This completes the proof.

Q.E.D.

Let us recall that the mapping class group $\Gamma_{S^1 \times S^1}$ of the torus $S^1 \times S^1$ is isomorphic to the group $SL_2(\mathbb{Z})$ of integral 2×2 -matrices with determinant

1. It is well-known that this group is generated by $S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and

$T = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ with relations $S^4 = I$, $(ST)^3 = S^2$. The matrices S and T correspond to the orientation preserving diffeomorphisms $S^1 \times S^1$ to $S^1 \times S^1$ which are defined by $S(z, w) = (\bar{w}, z)$ and $T(z, w) = (zw, w)$, $(z, w) \in S^1 \times S^1$, respectively, where we regard S^1 as the set of complex numbers of

absolute value 1. To define the Verlinde basis, we need one more orientation preserving diffeomorphism $U : S^1 \times S^1 \longrightarrow -S^1 \times S^1$ defined by $U(z, w) = (z, \bar{w})$ for $(z, w) \in S^1 \times S^1$ (see Figure 6).

Definition 2.5. *Let Z be a $(2+1)$ -dimensional TQFT. A basis $\{v_i\}_{i=0}^m$ of $Z(S^1 \times S^1)$ is said to be a Verlinde basis if it has the following properties.*

- (i) v_0 is the identity element of the fusion algebra associated with Z .
- (ii) (a) $Z(S)$ is presented by a unitary and symmetric matrix with respect to the basis $\{v_i\}_{i=0}^m$.
 (b) $Z(S)^2 v_0 = v_0$, and $Z(S)^2 v_i \in \{v_j\}_{j=0}^m$ for all i .
 (c) We define $S_{ji} \in \mathbb{C}$ by $Z(S)v_i = \sum_{j=0}^m S_{ji} v_j$. Then,
 1. $S_{i0} \neq 0$ for all i .
 2. $N_{ij}^k := \sum_{l=0}^m \frac{S_{il} S_{jl} \overline{S_{lk}}}{S_{0l}}$ ($i, j, k = 0, 1, \dots, m$) coincide with the structure constants of the fusion algebra with respect to $\{v_i\}_{i=0}^m$.
- (iii) $Z(T)$ is presented by a diagonal matrix with respect to the basis $\{v_i\}_{i=0}^m$.
- (iv) $Z(U)v_i = v_i^*$ for all i under the identification $Z(-S^1 \times S^1) \cong Z(S^1 \times S^1)^*$.

We call a pre-Verlinde basis a basis $\{v_i\}_{i=0}^m$ of $Z(S^1 \times S^1)$ satisfying the four conditions (i), (ii.a), (ii.b) and $Z(U)v_0 = v_0^*$.

Remarks 2.6. 1. If $\{v_i\}_{i=0}^m$ is a pre-Verlinde basis, then S_{i0} is a real number for all i .

2. A Verlinde basis is unique up to order of elements, since $w_i = S_{0i} \sum_{j=0}^m \overline{S_{ji}} v_j$ ($i = 0, 1, \dots, m$) are all orthogonal primitive idempotents in the fusion algebra satisfying $1 = w_0 + w_1 + \dots + w_m$. This fact follows from that the S -matrix diagonalizes the fusion rules in conformal field theory [23].

3. The map $\cdot : \{0, 1, \dots, m\} \longrightarrow \{0, 1, \dots, m\}$ defined by $Z(S)^2 v_i = v_i$ is an involution satisfying $\bar{0} = 0$.

4. The last condition (iv) was introduced in [24] and modified in [19].

Let us describe some basic results on a $(2+1)$ -dimensional TQFT with a pre-Verlinde basis.

Lemma 2.7. *Let Z be a $(2+1)$ -dimensional TQFT and $\{v_i\}_{i=0}^m$ a pre-Verlinde basis. Then, for the cobordism $W = (-D^2 \times S^1; S^1 \times S^1, \emptyset)$, we have $Z_W = v_0^* : Z(S^1 \times S^1) \longrightarrow \mathbb{C}$. In particular, $Z(S^2 \times S^1) = 1$.*




Figure 6: the action of $SL_2(\mathbb{Z})$

Proof. By considering the orientation preserving diffeomorphism $\tilde{U} : D^2 \times S^1 \longrightarrow -D^2 \times S^1$ defined by $\tilde{U}(z, w) = (z, \bar{w})$, we have $(Z(\tilde{U}|_{\partial D^2 \times S^1}))(Z(D^2 \times S^1)) = Z(-D^2 \times S^1) \in Z(-S^1 \times S^1)$. Since $\tilde{U}|_{\partial D^2 \times S^1} = U$ and $Z(D^2 \times S^1) = v_0$, it follows that $v_0^* = Z(-D^2 \times S^1) = Z_W$ as elements in $Z(S^1 \times S^1)^*$.

Next, we prove that $Z(S^2 \times S^1) = 1$. The 3-manifold $S^2 \times S^1$ is regarded as $(D_+^2 \times S^1) \cup (-D_-^2 \times S^1)$, where $D_+^2 = \{(x, y, z) \in S^2 \mid z \geq 0\}$, $D_-^2 = \{(x, y, z) \in S^2 \mid z \leq 0\}$. Thus, for the cobordisms $W_1 := (D_+^2 \times S^1; \emptyset, \partial D_+^2 \times S^1)$ and $W_2 := (-D_-^2 \times S^1; \partial D_-^2 \times S^1, \emptyset)$, we have

$$Z(S^2 \times S^1) = Z_{W_2} \circ Z_{W_1}.$$

Since $W_1 \cong (D^2 \times S^1; \emptyset, S^1 \times S^1) = W_0$ and $W_2 \cong (-D^2 \times S^1; S^1 \times S^1, \emptyset) = W$, it follows that

$$Z(S^2 \times S^1) = Z_W \circ Z_{W_0} = v_0^*(v_0) = 1.$$

This completes the proof.

Q.E.D.

Lemma 2.8. *Let Z be a $(2+1)$ -dimensional TQFT and $\{v_i\}_{i=0}^m$ a pre-Verlinde basis. Then, $J(\bigcirc; i) = S_{i0}$ for each $i \in \{0, 1, \dots, m\}$.*

Proof. Let K be the trivial knot with 0-framing given by unit circle in $\mathbb{R}^3$, and $X = \overline{S^3 - N(K)}$ the knot exterior of K . The cobordism $W_K = (X; \partial N(K), \emptyset)$ is isomorphic to the cobordism $W := (-D^2 \times S^1; S^1 \times S^1, \emptyset)$ via the orientation preserving diffeomorphism $f : X \longrightarrow -D^2 \times S^1$ such that $f(M) = l$ and $f(L) = -m$, where M and L are simple closed curves on $\partial N(K)$ depicted in Figure 7.

Then, we have

$$Z_{W_K} = Z_W \circ Z(f|_{\partial N(K)}).$$

Since

$$Z(f|_{\partial N(K)}) \circ Z(j) = Z(S)$$




Figure 7:

for $j := h|_{S^1 \times S^1}$, where $h : D^2 \times S^1 \longrightarrow N(K)$ is the framing of K , we have

$$J(K, i) = \langle v_0^*, Z(S)(v_i) \rangle = S_{i0}.$$

This completes the proof.

Q.E.D.

The framed link invariants $J(L; i_1, \dots, i_r)$, $(i_1, \dots, i_r = 0, 1, \dots, m)$ have the following nice properties.




Figure 8: a positive curl and a negative curl

Lemma 2.9. *Let Z be a $(2 + 1)$ -dimensional TQFT and $\{v_i\}_{i=0}^m$ a pre-Verlinde basis. For a framed link $L = L_1 \cup L_2 \cup \dots \cup L_r$, the invariant $J(L; i_1, \dots, i_r)$ has the following properties.*

(1) *Let $L' = L_1 \cup \dots \cup (-L_k) \cup \dots \cup L_r$ be the framed link obtained from L by changing the orientation for the k -th component L_k . Then,*

$$J(L; i_1, \dots, i_k, \dots, i_r) = J(L'; i_1, \dots, \bar{i}_k, \dots, i_r).$$

(2) *Suppose that $\{v_i\}_{i=0}^m$ satisfies the condition (iii) in the definition of Verlinde basis. We define $t_i \in \mathbb{C}$ ($i = 0, 1, \dots, r$) by $Z(T)v_i = t_i v_i$. Let L' be the framed link obtained from L such that it is same as L except for a small segment of the k -th component L_k and the small segment is replaced by a*

positive or negative curl as shown in Figure 8. Then, the following equations hold.

If the small segment is replaced by a positive curl, then

$$J(L'; i_1, \dots, i_r) = t_{i_k}^{-1} J(L; i_1, \dots, i_r).$$

If the small segment is replaced by a negative curl, then

$$J(L'; i_1, \dots, i_r) = t_{i_k} J(L; i_1, \dots, i_r).$$

Proof. Without loss of generality, we may suppose that $k = 1$.

(1) Let $h_i : D^2 \times S^1 \rightarrow N(L_i)$ be the framing of L_i for each $i = 1, \dots, r$, and $h'_1 : D^2 \times S^1 \rightarrow N(-L_1)$ the framing of $-L_1$. We set $j_i := h_i|_{S^1 \times S^1}$ for each $i = 1, \dots, r$, and $j'_1 := h'_1|_{S^1 \times S^1}$. Then,

$$j'_1 = j_1 \circ S^2,$$

since $h'_1 = h_1 \circ f$ for the orientation preserving diffeomorphism $f : D^2 \times S^1 \rightarrow D^2 \times S^1$ defined by $f(z, w) = (\bar{z}, \bar{w})$ (see Figure 9).




Figure 9:

Let X be the link exterior of L . Since the two cobordisms

$$\mathcal{W}_L = (X; \prod_{i=1}^r j_i, \emptyset), \quad \mathcal{W}_{L'} = (X; j'_1 \prod_{i=2}^r j_i, \emptyset)$$

are isomorphic via the identity map id_X , we see that

$$Z_{\mathcal{W}_{L'}} \circ (Z(S^2) \circ id \otimes \dots \otimes id) = Z_{\mathcal{W}_L}.$$

Since $Z(S^2)v_i = v_{\bar{i}}$, we have

$$J(L; i_1, i_2, \dots, i_r) = J(L'; \bar{i}_1, i_2, \dots, i_r).$$

(2) We suppose that L' arises from L by replacing a small segment of the first component L_1 by a positive curl. Let $h'_1 : D^2 \times S^1 \rightarrow N(L'_1)$ be the framing of L'_1 . We set $j'_1 := h'_1|_{S^1 \times S^1}$.

Let X' be the link exterior of L' . Then, there exists an orientation preserving diffeomorphism $f : X' \rightarrow X$ such that




Figure 10:

- (i) $j_1^{-1} \circ (-f|_{\partial N(L_1)}) \circ j'_1 = T^{-1}$ up to isotopy,
- (ii) $-f|_{\partial N(L_i)} = id_{\partial N(L_i)}$ for all $i = 2, \dots, r$.

This map f gives rise to the isomorphism between the cobordisms $\mathcal{W}_{L'} = (X'; \emptyset, j'_1 \amalg (\amalg_{i=2}^r j_i))$ and $\mathcal{W}_L = (X; \emptyset, \amalg_{i=1}^r j_i)$. Hence, we have

$$Z_{\mathcal{W}_{L'}} \circ (Z(j_1'^{-1}) \otimes (\bigotimes_{i=2}^r Z(j_i^{-1}))) = Z_{\mathcal{W}_L} \circ (\bigotimes_{i=1}^r Z(j_i^{-1}) \circ Z(-f|_{\partial N(L_i)})).$$

It follows that

$$Z_{\mathcal{W}_{L'}} = Z_{\mathcal{W}_L} \circ (Z(T^{-1}) \otimes id \cdots \otimes id).$$

This implies that

$$J(L'; i_1, \dots, i_r) = t_{i_1}^{-1} J(L; i_1, \dots, i_r).$$

Thus, the proof of the first equation of (2) is completed. By a similar argument, the second equation of (2) can be proved. This completes the proof. Q.E.D.

Let us introduce a criterion for the Verlinde identity (ii.c.2).

We suppose that a $(2+1)$ -dimensional TQFT Z arises from a semisimple ribbon $\mathbb{C}$ -linear Ab-category in the sense of Turaev [21]. Let $\{v_i\}_{i=0}^m$ be a pre-Verlinde basis satisfying the equivalent conditions (1) and (2) in Proposition 2.4. We suppose $S_{i0} \neq 0$ for all $i = 0, 1, \dots, m$.

For the 3-component framed link L presented by the diagram as in Figure 11, the framed link invariant $J(L; i, j, k)$ coincides with the quantum trace of $J(T_{i,l}) \circ J(T_{j,l})$, where $T_{i,l}$ is the colored framed tangle presented by the diagram as in Figure 12.




Figure 11:

Since $J(T_{i,l})$ is a map from the simple object l to l , the invariant $J(T_{i,l})$ is a scalar multiple by id_l . We define a_i to be this scalar. Since the quantum trace of $T_{i,l}$ is the Hopf link with colors i, l , we see that

$$a_i \cdot J(\bigcirc; l) = S_{il}.$$

This implies that

$$J(L; i, j, l) = \frac{S_{il}S_{jl}}{S_{l0}}. \quad (2.1)$$




Figure 12: the colored 1 – 1 tangle $T_{i,l}$

On the other hand, by fusing L_i and L_j we have

$$J(L; i, j, l) = \sum_{i=0}^m N_{ij}^k S_{kl}, \quad (2.2)$$

where N_{ij}^k ($i, j, k = 0, 1, \dots, m$) are the structure constants of the fusion algebra with respect to $\{v_i\}_{i=0}^m$. From (2.1) and (2.2), it follows that

$$\sum_{i=0}^m N_{ij}^k S_{kl} = \frac{S_{il}S_{jl}}{S_{l0}}.$$

Since the above equation induces the Verlinde identity

$$N_{ij}^k = \sum_{l=0}^m \frac{S_{il} S_{jl} \overline{S_{lk}}}{S_{l0}},$$

the condition (2) in Proposition 2.4 implies the condition (ii.c.2) in the definition of Verlinde basis (see [20], [25] for similar arguments). If $\{v_i\}_{i=0}^m$ satisfies the condition (iii) in the definition of Verlinde basis, then we see that the converse is true by using the technique in [11] (see Figure 13).




Figure 13:

Thus, we conclude the following.

Theorem 2.10. *Suppose that a $(2+1)$ -dimensional TQFT Z arises from a semisimple ribbon $\mathbb{C}$ -linear Ab-category. Let $\{v_i\}_{i=0}^m$ be a pre-Verlinde basis satisfying $S_{i0} \neq 0$ for all $i = 0, 1, \dots, m$. If $Z(U)v_i = v_i^*$ for all $i = 0, 1, \dots, m$, then $N_{ij}^k := \sum_{l=0}^m \frac{S_{il} S_{jl} \overline{S_{lk}}}{S_{l0}}$ ($i, j, k = 0, 1, \dots, m$) coincide with the structure constants of the fusion algebra with respect to $\{v_i\}_{i=0}^m$.*

Furthermore, if $\{v_i\}_{i=0}^m$ satisfies the condition (iii) in the definition of Verlinde basis, then the converse is true. Therefore, the following are equivalent.

- (i) $J(H; i, j) = S_{ij}$ for all $i, j = 0, 1, \dots, m$.
- (ii) $Z(U)v_i = v_i^*$ for all $i = 0, 1, \dots, m$.
- (iii) $N_{ij}^k := \sum_{l=0}^m \frac{S_{il} S_{jl} \overline{S_{lk}}}{S_{l0}}$ ($i, j, k = 0, 1, \dots, m$) coincide with the structure constants of the fusion algebra with respect to $\{v_i\}_{i=0}^m$.

Here, H is the Hopf link presented by the diagram as in Figure 3.

3 Turaev-Viro-Ocneanu $(2+1)$ -dimensional TQFT (Review)

3.1 Sectors and finite system Δ

For a detailed exposition about sectors, see [6]. Let $N \subset M$ be an inclusion of infinite factors. In this case, we also have a similar concept to Jones

index named Kosaki index, and if it takes the minimal value, we write it $[M : N]_0$ as in the case of Jones index. In the sequel, we assume subfactors have finite minimal indices. For the inclusion of $\rho(M) \subset M$, where $\rho \in \text{End}(M)$, we call $[M : \rho(M)]_0^{1/2}$ the statistical dimension of ρ and denote it by $d(\rho)$. We denote the set of $*$ -homomorphisms from N to M with finite statistical dimensions by $\text{Mor}(N, M)_0$. We say that $\rho_1, \rho_2 \in \text{Mor}(N, M)_0$ are equivalent if there exists a unitary u in M such that $u\rho_1(x) = \rho_2(x)u$ for all $x \in N$. This gives an equivalence relation in $\text{Mor}(N, M)_0$. The set of the equivalence classes is denoted by $\text{Sect}(N, M)$ and its element is called an M - N sector, which is denoted by $[\rho]$ for $\rho \in \text{Mor}(N, M)_0$. In M - M sectors $\text{Sect}(M, M) =: \text{Sect}(M)$, we have the product $[\rho_1] \cdot [\rho_2] = [\rho_1 \circ \rho_2]$ and the summation $[\rho_1] \oplus [\rho_2]$. (See [6] for the definition of the summation of sectors.) These operations define a semiring structure in $\text{Sect}(M)$. For $[\rho_1], [\rho_2] \in \text{Sect}(M)$, we define

$$(\rho_1, \rho_2) = \{V \in M \mid V\rho_1(x) = \rho_2(x)V, \forall x \in M\}.$$

It is called the *intertwiner space* between ρ_1 and ρ_2 . If $(\rho, \rho) = \mathbb{C}1_M$, we say that ρ is *irreducible*. The intertwiner space (ρ_1, ρ_2) has an inner product $\langle V, W \rangle = W^* \cdot V$, $V, W \in (\rho_1, \rho_2)$, if ρ_1 is irreducible.

Moreover, $\text{Sect}(M)$ has a conjugation $\overline{[\rho]} = [\bar{\rho}]$. Namely, for $[\rho] \in \text{Sect}(M)$, there exist an endomorphism $\bar{\rho} \in \text{End}(M)$ and a pair of intertwiners $R_\rho \in (id, \bar{\rho})$ and $\bar{R}_\rho \in (id, \rho\bar{\rho})$ such that $\bar{R}_\rho^* R_\rho = R_\rho^* \bar{\rho}(\bar{R}_\rho) = 1/d(\rho)$. With these operations, $\text{Sect}(M)$ becomes a $*$ -semiring over $\mathbb{C}$.

An important thing is that $\text{Sect}(M)$ is closed under the operations such as product, direct sum, irreducible decomposition and conjugation.

Let us introduce the notion of a finite system Δ of $\text{End}(M)_0$, which is a basic data to describe a topological quantum field theory from subfactors. Since the embedding $\iota : N \hookrightarrow M$ is an element of $\text{Mor}(N, M)_0$, we can consider the sector $[\iota] \in \text{Sect}(N, M)$. We note that the conjugation $\overline{[\iota]}$ is an element of $\text{Sect}(M, N)$, and the product $[\iota]\overline{[\iota]}$ becomes an element of $\text{Sect}(M)$. In a similar way, $\overline{[\iota]}[\iota]$ becomes an element of $\text{Sect}(N)$. By decomposing $([\iota]\overline{[\iota]})^n$, $(\overline{[\iota]}[\iota])^n$, $[\iota](\overline{[\iota]}[\iota])^n$ and $(\overline{[\iota]}[\iota])^n$ into irreducible sectors, we get M - M , M - N , N - M and N - N sectors responsibly. (A sector $[\rho]$ is said to be *irreducible* if ρ is irreducible.) If the number of the irreducible sectors in the above decompositions is finite, then the subfactor $N \subset M$ is called of *finite depth*. Throughout this paper, we only consider subfactors with finite depth and finite index. For a finite depth subfactor, we get finitely many irreducible M - M sectors. In other words, we have a representative set $\Delta = \{\rho_\xi\}_{\xi \in \Delta_0}$ of finitely many irreducible M - M sectors such that

- (i) $[\rho_\xi] = [\rho_\eta]$ if and only if $\xi = \eta$
- (ii) There exists $e \in \Delta_0$ such that $\rho_e = id$
- (iii) For any $\xi \in \Delta_0$, there exists $\bar{\xi} \in \Delta_0$ such that $\overline{[\rho_\xi]} = [\rho_{\bar{\xi}}]$
- (iv) There exist non-negative integers $N_{\xi, \eta}^\zeta$ such that $[\rho_\xi][\rho_\eta] = \bigoplus_{\zeta \in \Delta_0} N_{\xi, \eta}^\zeta [\rho_\zeta]$.

We call this Δ a finite system of $\text{End}(M)_0$ or simply a *finite system*.

Remark 3.1. We note that Δ can be seen as a C^* -tensor category in the following manner. The objects of the category are $\mathbb{C}$ -linear span of elements of Δ , the morphisms of the category are intertwiners, and the tensor product structure is given by the compositions of endomorphisms in objects of the category. By an abuse of notation, we denote this category by Δ . See [12] for the details.

3.2 Turaev-Viro-Ocneanu TQFT

We need some preparations before constructing the Turaev-Viro-Ocneanu TQFT. In the sequel, we write ξ instead of ρ_ξ and so forth, for simplicity.

Let Δ be the finite system of $\text{End}(M)_0$. We consider the following diagram.

$$\begin{array}{ccc} \xi \cdot \alpha \cdot \eta & \xleftarrow{\xi(T_1)} & \xi \cdot \beta \\ T_3 \uparrow & & \uparrow T_2 \\ \gamma \cdot \eta & \xleftarrow{T_4} & \delta \end{array}$$

Here, $\alpha, \beta, \gamma, \xi, \eta, \delta \in \Delta$ and $T_1 \in (\beta, \alpha \cdot \eta)$, $T_2 \in (\delta, \xi \cdot \beta)$, $T_3 \in (\gamma, \xi \cdot \alpha)$, $T_4 \in (\delta, \gamma \cdot \eta)$. Then, the composition of intertwiners $T_4^* \cdot T_3^* \cdot \xi(T_1) \cdot T_2$ belongs to (δ, δ) . Since δ is assumed to be irreducible, this composition of intertwiners is regarded as a complex number. We call this number a *quantum 6j-symbol*.

The above diagram can be seen as a tetrahedron as in Figure 14.




Figure 14: A tetrahedron as a diagram

We assign $d(\beta)^{-1/2}d(\gamma)^{-1/2}T_4^* \cdot T_3^* \cdot \xi(T_1) \cdot T_2$ to this tetrahedron.

Let V be an oriented closed 3-dimensional manifold. Choose a triangulation of V and write it $\mathcal{T}$. To each vertex in $\mathcal{T}$, we assign the factor M , to each edge in $\mathcal{T}$ an element in Δ and to each face in $\mathcal{T}$ an intertwiner. Let E be the set of the edges in $\mathcal{T}$, e be an assignment of elements in Δ to the edges in $\mathcal{T}$, and φ be an assignment of intertwiners to the faces in $\mathcal{T}$. A tetrahedron τ has the labeled edges by the assignment e and the labeled faces by the assignment φ . Hence, to a tetrahedron τ , we assign a complex number defined by using the 6j-symbol as above or its complex conjugate depending on the orientation of τ . We denote this complex number by $W(\tau; e, \varphi)$. We multiply

these W 's and the weights $\prod_E d(\xi)$. Then, sum up all these resulting values. Finally, we multiply some weights coming from the vertices in $\mathcal{T}$.

$$Z_\Delta(V, \mathcal{T}) = \lambda^{-a} \sum_e \left(\prod_E d(\xi) \right) \sum_\varphi \prod_\tau W(\tau; e, \varphi),$$

where $\lambda = \sum_{\xi \in \Delta_0} d(\xi)^2$ and a is the number of the vertices in $\mathcal{T}$. This $Z_\Delta(V, \mathcal{T})$ is proven to be independent of any choice of triangulations because of the properties of quantum $6j$ -symbols. (See [2] for a detailed account.) Namely, $Z_\Delta(V, \mathcal{T})$ turns out to be a topological invariant of V . So, we drop $\mathcal{T}$ off from $Z_\Delta(V, \mathcal{T})$ and denote this value by $Z_\Delta(V)$. We call $Z_\Delta(V)$ the *Turaev-Viro-Ocneanu invariant* of the 3-dimensional manifold V . When V is an oriented, compact 3-dimensional manifold possibly with boundary, first we fix a triangulation $\partial\mathcal{T}$ of the boundary of V and extend it to the whole triangulation of V . Then, we assign an element in Δ to each edge in $\partial\mathcal{T}$ and assign an intertwiner to each face. We fix these assignments to the end and denote them by ∂e and $\partial\varphi$, respectively. In a similar way to the closed case, we assign M , an element in Δ and an intertwiner to each vertex, each edge and each face in the triangulation $\mathcal{T} \setminus \partial\mathcal{T}$. We make $Z_\Delta(V, \mathcal{T}, \partial e, \partial\varphi)$ as above:

$$Z_\Delta(V, \mathcal{T}, \partial e, \partial\varphi) = \lambda^{-a+\partial a/2} \prod_{\partial e} d(\xi)^{1/2} \sum_{e \setminus \partial e} \left(\prod_{E \setminus \partial E} d(\xi) \right) \sum_\varphi \prod_\tau W(\tau; e, \varphi),$$

where ∂a is the number of the vertices on the boundary. This value $Z_\Delta(V, \mathcal{T}, \partial e, \partial\varphi)$ does not depend on the assignments of the vertices, the edges and the faces in $\mathcal{T} \setminus \partial\mathcal{T}$. Such extended Z_Δ gives rise to a unitary TQFT because to an oriented closed surface, it assigns a finite dimensional Hilbert space with the inner product induced from the space of intertwiners. (See [2] for the detailed construction.) We call this TQFT the $(2+1)$ -dimensional *Turaev-Viro-Ocneanu TQFT* and denote it by Z_Δ , again.

4 Verlinde basis of $Z_\Delta(S^1 \times S^1)$

Let $N \subset M$ be a subfactor of an infinite factor M with finite index and finite depth, and let Δ be a finite system of irreducible M - M endomorphisms arising from the subfactor. We write, for instance, ξ instead of ρ_ξ and so forth for the elements of Δ .

Based on Δ , we construct a new finite dimensional C^* -algebra named the *tube algebra* $\text{Tube}\Delta$ as in [14]. (Also see [7], but our normalization convention is different from that there.) In this section, $\text{Tube}\Delta$ plays a crucial role to find a nicely behaved basis of $Z_\Delta(S^1 \times S^1)$, which we call a Verlinde basis. It makes the Turaev-Viro-Ocneanu TQFT a rich theory.

We sometimes use the simple notation $Z(S^1 \times S^1)$ instead of $Z_\Delta(S^1 \times S^1)$ in the sequel.

4.1 Tube algebras

A tube algebra $\text{Tube}\Delta$, which was first introduced by Ocneanu in [14], is defined by $\bigoplus_{\xi, \eta, \zeta} (\xi \cdot \zeta, \zeta \cdot \eta)$ as a vector space over $\mathbb{C}$. Its element is a linear combination of the composition $T_2 \cdot T_1^*$ of the orthonormal bases of intertwiner spaces $T_1 \in (\delta, \xi \cdot \zeta)$, $T_2 \in (\delta, \zeta \cdot \eta)$, where ξ, η, ζ and δ run over Δ_0 . An element in $\text{Tube}\Delta$ can be depicted as in the left-hand side of Figure 16. We will define a product structure and a $*$ -structure on it.

The product structure

Let $X = X_2 X_1^* \in (\xi \cdot \zeta, \zeta \cdot \eta)$ for $X_1 \in (\delta, \xi \cdot \zeta)$, $X_2 \in (\delta, \zeta \cdot \eta)$ and $Y = Y_2 Y_1^* \in (\xi' \cdot \zeta', \zeta' \cdot \eta')$ for $Y_1 \in (\delta', \xi' \cdot \zeta')$, $Y_2 \in (\delta', \zeta' \cdot \eta')$. Then, the product of X and Y in $\text{Tube}\Delta$ is defined by the following formula.

$$X \cdot Y = \delta_{\eta, \xi'} \lambda \sum d(\xi)^{-1/2} d(\eta')^{-1/2} d(\eta) \lambda(X, Y; Z) Z.$$

Here, the summation is taken over $Z = Z_2 Z_1^*$, $Z_1 \in (\tau, \xi \cdot \nu)$, $Z_2 \in (\tau, \nu \cdot \eta')$, where Z_1 and Z_2 are orthonormal bases of the intertwiner spaces $(\tau, \xi \cdot \nu)$ and $(\tau, \nu \cdot \eta')$ respectively, and $\lambda(X, Y; Z)$ is the value of Z_Δ of the 3-manifold depicted in Figure 15. For general elements X, Y in $\text{Tube}\Delta$, we define the product of them by linearity since X and Y are linear combinations of the forms $X_2 X_1^*$ and $Y_2 Y_1^*$ as above.




Figure 15: A coefficient of the product of $X \cdot Y$

The $*$ -structure

Let X be as above. Then, we can consider X as a tube as in the right-hand side of Figure 16. The $*$ -operation is defined by the inverting the tubes inside out. We denote this $*$ -operation by X^* . See Figure 17.




Figure 16: An element of the tube algebra as a tube




Figure 17: An inversed tube

With the product and the $*$ -structure defined as above, $\text{Tube}\Delta$ becomes a finite dimensional C^* -algebra. Since any finite dimensional C^* -algebra is semisimple, we may assume that $\text{Tube}\Delta \cong \bigoplus_{i=0}^r M_{n_i}(\mathbb{C})$.

We observe that the definition of tube algebras is compatible with the operations of Turaev-Viro-Oceanu TQFT such as gluing, cutting and so forth.

Remark 4.1. In [7], M. Izumi has introduced the tube algebra in the setting of sectors, but it is slightly different from ours in the normalization coefficients. We followed the definition of the tube algebra in [2, Chapter 12].

Before we start the analysis of tube algebras, we list the notations that we will use frequently.

For the solid torus $D^2 \times S^1$, we denote the value of Z_Δ of $D^2 \times S^1$ assigned a vector λ on the boundary by $Z_\Delta(D^2 \times S^1; \lambda)$.

For the 3-manifold $D^2 \times S^1 \setminus \text{Int}D_0^2 \times S^1$, where D_0 is contained in $\text{Int}D^2$, we denote the value of Z_Δ of $D^2 \times S^1 \setminus \text{Int}D_0^2 \times S^1$ assigned vectors λ and μ , λ for the boundary of $D_0^2 \times S^1$ and μ for the boundary of $D^2 \times S^1$ by $Z_\Delta(D^2 \times S^1 \setminus \text{Int}D_0^2 \times S^1; \lambda, \mu)$. See Figure 18.




Figure 18: The labeled 3-manifold obtained by removing another solid torus from a solid torus

In a similar manner, for the 3-manifold $D^2 \times S^1 \setminus (\text{Int}D_1^2 \times S^1 \cup \text{Int}D_2^2 \times S^1)$, where D_1^2 and D_2^2 are two disjoint disks contained in $\text{Int}D^2$, we denote the value of Z_Δ of $D^2 \times S^1 \setminus (\text{Int}D_1^2 \times S^1 \cup \text{Int}D_2^2 \times S^1)$ assigned vectors λ , μ and ν , λ for the boundary of $D^2 \times S^1$, μ for the boundary of $D_1^2 \times S^1$ and ν for the boundary of $D_2^2 \times S^1$, by $Z_\Delta(D^2 \times S^1 \setminus (\text{Int}D_1^2 \times S^1 \cup \text{Int}D_2^2 \times S^1); \lambda, \mu, \nu)$. See Figure 19.

Let $\{\pi_0, \dots, \pi_r\}$ be the minimal central projections of $\text{Tube}\Delta$. Then, we have the following lemma.




Figure 19: The labeled 3-manifold obtained by removing two solid tori from a solid torus

Lemma 4.2. $\langle \pi_i, \pi_j \rangle_{Z(S^1 \times S^1)} = \delta_{i,j} n_i^2$ ($i, j = 0, \dots, r$), where $\langle \cdot, \cdot \rangle_{Z(S^1 \times S^1)}$ is the inner product of $Z_\Delta(S^1 \times S^1)$ defined by the Turaev-Viro-Ocneanu TQFT.

Proof. First, we consider the case $i = j$. We note that $\langle \pi_i, \pi_i \rangle_{Z(S^1 \times S^1)} = Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; \pi_i, \pi_i)$.

Cut $D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1$ along the meridian, then $T = \text{annulus} \times [0, 1]$ is created. Let A_{jk} be the value of $Z_\Delta(T, \xi_j, \xi_k)$, where T is labeled by ξ_j on one side of the sections and by ξ_k on the other side. Then, $Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; \pi_i, \pi_i) = \sum_{\xi_j} A_{jj}$. There exists an operator A such that $A_{jk} = \langle A\xi_j, \xi_k \rangle$, where $\langle \cdot, \cdot \rangle$ is the inner product on intertwiners on the sections of T so that $\langle \xi_j, \xi_k \rangle = \delta_{jk}$.

It is easy to see that the operator A is a projection. Hence, $\langle \pi_i, \pi_i \rangle_{Z(S^1 \times S^1)} = \sum_{j=1}^m A_{jj} = \text{Tr}(A) \in \mathbb{N}$. Namely, we have proved that $\langle \pi_i, \pi_i \rangle_{Z(S^1 \times S^1)} = \dim \pi_i(\text{Tube} \Delta) = n_i^2$.

When $i \neq j$, it is easy to see that $\langle \pi_i, \pi_j \rangle_{Z(S^1 \times S^1)} = 0$, because π_i and π_j are central projections orthogonal to each other. This ends the proof. Q.E.D.

Theorem 4.3. Let Δ be a finite system. Then, the center of $\text{Tube} \Delta$ is naturally isomorphic to $Z_\Delta(S^1 \times S^1)$ as a vector space.

Proof. Let $V(S^1 \times S^1)$ be the linear span of elements in $\text{Tube} \Delta$ such that the left and right labels in Figure 16 are equal. Since $\text{Tube} \Delta \supset V(S^1 \times S^1)$ and $V(S^1 \times S^1)$ contains the center of $\text{Tube} \Delta$, it is enough to consider $V(S^1 \times S^1)$ instead of whole tube algebra.

For $X \in V(S^1 \times S^1)$, we set $\varphi_i(X) = Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; \pi_i, X)$. If $X \perp V(S^1 \times S^1)$ with respect to the colored inner product of $\text{Tube} \Delta$, we set $\varphi_i(X) = 0$. Here, for $X = X_2 X_1^* \in (\xi \cdot \zeta, \zeta \cdot \eta)$, $X_1 \in (\delta, \xi \cdot \zeta)$, $X_2 \in (\delta, \zeta \cdot \eta)$ and $Y = Y_2 Y_1^* \in (\xi' \cdot \zeta', \zeta' \cdot \eta')$, $Y_1 \in (\delta', \xi' \cdot \zeta')$, $Y_2 \in (\delta', \zeta' \cdot \eta')$, the colored inner product $\langle X, Y \rangle_{\text{color}}$ of $\text{Tube} \Delta$ is defined by $\delta_{\xi, \xi'} \delta_{\eta, \eta'} \delta_{\zeta, \zeta'} \langle X_1, Y_1 \rangle \langle X_2, Y_2 \rangle$. The last two brackets stand for the inner products of the intertwiner spaces $(\delta, \xi \cdot \zeta)$ and $(\delta, \zeta \cdot \eta)$, respectively. Then, this φ_i is a linear functional defined on $\text{Tube} \Delta$. It is easy to see that φ_i is tracial.

If X is an element in $\pi_i(\text{Tube} \Delta)$, then we have

$$\varphi_i(X) = \langle \pi_i, \pi_i \rangle_{Z(S^1 \times S^1)} \text{tr}(X),$$

where tr is the normalized trace. In a similar way, for $X, Y \in \pi_i(\text{Tube}\Delta)$, the value $Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; Y, X)$ is equal to $\langle \pi_i, \pi_i \rangle_{Z(S^1 \times S^1)} \text{tr}(X) \text{tr}(Y^*)$.

When we write $X_j = \pi_j X$ for $X \in V(S^1 \times S^1)$, $X = \bigoplus_{j=0}^r X_j$. For $X \in \text{Tube}\Delta$, we put

$$E(X) = \sum_{i=0}^r \frac{\varphi_i(X)}{\langle \pi_i, \pi_i \rangle_{Z(S^1 \times S^1)}} \pi_i = \sum_{i=0}^r \text{tr}(X_i) \pi_i.$$

Then, this is a conditional expectation from $\text{Tube}\Delta$ to the center of $\text{Tube}\Delta$. We have a description of the kernel of E by $\text{Ker} E = \text{the linear span of } \{X \in V(S^1 \times S^1) \mid \text{tr}(X_j) = 0, j = 0, \dots, r\} \cup \{X \in \text{Tube}\Delta \mid \langle X, Y \rangle_{\text{color}} = 0 \text{ for all } Y \in V(S^1 \times S^1)\}$.

We have

$$\begin{aligned} \langle X_j, X_j \rangle_{Z(S^1 \times S^1)} &= \langle \pi_j, \pi_j \rangle_{Z(S^1 \times S^1)} \text{tr}(X_j) \text{tr}(X_j^*) \\ &= \langle \pi_j, \pi_j \rangle_{Z(S^1 \times S^1)} |\text{tr}(X_j)|^2, \end{aligned}$$

and we put $Q = \{X \in V(S^1 \times S^1) \mid \langle X, X \rangle_{Z(S^1 \times S^1)} = 0\}$. Then, $Z(S^1 \times S^1) = V(S^1 \times S^1)/Q = \text{Tube}\Delta/\text{Ker} E \cong \text{Center}(\text{Tube}\Delta)$. Precisely, denoting the embedding map from $Z_\Delta(S^1 \times S^1)$ into $\text{Tube}\Delta$ by ι , we have proved that $\text{Center}(\text{Tube}\Delta) = \iota(Z_\Delta(S^1 \times S^1)) \subset \text{Tube}\Delta$. Q.E.D.

Remark 4.4. *From this theorem and Lemma 4.2, $\{\frac{\pi_i}{n_i}\}_{i=0}^r$ is an orthonormal basis of $Z(S^1 \times S^1)$.*

4.2 Verlinde basis

In this subsection, we show the existence of a basis of $Z_\Delta(S^1 \times S^1)$ nicely behaved under the action of $SL_2(\mathbb{Z})$, which we call the *Verlinde basis*. It deeply depends on the structure of the tube algebra.

Before the proof of the existence of such basis, we need some preparations.

Let p_i be a minimal projection in $\pi_i(\text{Tube}\Delta)$ for each $i = 0, \dots, r$. By Lemma 4.2, we have $\langle p_i, p_j \rangle_{Z(S^1 \times S^1)} = \delta_{ij}$. (We use the same notation p_i as an element of $Z_\Delta(S^1 \times S^1)$.) Then, by the last remark in the previous subsection, we have

$$p_i = \frac{\pi_i}{n_i} \quad (i = 0, \dots, r) \tag{4.3}$$

in $Z_\Delta(S^1 \times S^1)$.

Let us compute the value $Z_\Delta(D^2 \times S^1; p_i)$. For this, we look at $Z_\Delta(D^2 \times S^1; \pi_i)$. It is a summation of $Z_\Delta(D^2 \times [0, 1], \xi_k, \xi_k)$ over the intertwiners ξ_k 's since $D^2 \times [0, 1]$ is created by cutting the solid torus along the meridian. This is nothing but the summation of the dimension of the partially labeled disks labeled by p_{i_j} over $j, j = 1, \dots, n_i$, where p_{i_j} 's are minimal projections such that $\sum_j p_{i_j} = \pi_i$. (See [2, Chapter 12] for the definition of the partially labeled surfaces.) See Figure 20.




Figure 20: The decomposition of a labeled disk

When we denote the dimension of the partially labeled disk with label p_{i_j} by c_i , the dimension of the partially labeled disk with label π_i becomes $n_i c_i$.

Take a summation of $n_i c_i$ over all i 's, then it is equal to the dimension of the partially labeled disk with label $\sum_i \pi_i = 1$. Hence, it is equal to the dimension of the triangulated disk in Figure 21, which edge AB is glued together. (The boundary element is a direct sum taken over arbitray ρ .)




Figure 21: A triangulated disk

Then, this vector space is non-trivial only in the case of $\rho = id$ and then, the dimension is one. It means that $\sum_i n_i c_i = 1$. Hence, only one summand can survive. We may and do assume $n_0 c_0 = 1$. (Hence, $n_0 = 1$, $c_0 = 1$.) In a similar manner, we denote π_i by π_0 in this case.

Let us summerize the above argument as a lemma.

Lemma 4.5. $Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; \pi_i) = Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; p_i) = \delta_{i0}$.

Let N_{ij}^k be $Z_\Delta(D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1); p_i, p_j; p_k)$. See Figure 19. Cut $D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1)$ along the meridian, then we have $P = (D^2 \setminus \text{Int} D_1^2 \cup \text{Int} D_2^2) \times [0, 1]$, and as a value, N_{ij}^k is a summation of $Z_\Delta(P, \xi_k, \xi_k)$ over ξ_k 's, where P is labeled by ξ_k 's on the sections. This value can be written $\langle A \xi_k, \xi_k \rangle$, where $\langle \cdot, \cdot \rangle$ is the inner product of the Hilbert space of the section. Then, it is easy to see that A is a projection. Hence, $N_{ij}^k = \text{Tr}(A) = \dim H_{\text{pants}} \in \mathbb{N}$, where H_{pants} is the Hilbert space associated with the 3-holed sphere in the Turaev-Viro-Oceanu TQFT. See Figure 22.

With these settings, we can prove the following theorem, which is one of our main theorems in this paper. (Many of the contents have already appeared in [2], [3] following several presentations of Oceanu, but some normalizations are missing or incorrect there, so we include a complete proof here.)




Figure 22: H_{pants}

Theorem 4.6. *Let Δ be a finite system and $\{p_i\}_{i=0}^r$ minimal projections such that each p_i belongs to $\pi_i(\text{Tube}(\Delta))$, $i = 0, \dots, r$. Then, $\{p_i\}_{i=0}^r$ is a Verlinde basis of $Z_\Delta(S^1 \times S^1)$ in the sense of Section 2.*

Proof. We check all the conditions for $\{p_0, \dots, p_r\}$ to be a Verlinde basis step by step.

Condition (ii.c.1): Look at the inner product $n_j^2 = \langle \pi_j, \pi_j \rangle_{Z(S^1 \times S^1)}$. This value is defined by $Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; \pi_j, \pi_j)$. Cut $D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1$ along the meridian, then we have $X_0 = \text{annulus} \times [0, 1]$ as topological object. Since π_j is central, X_0 can be depicted as in Figure 23, where ξ_k 's are intertwiners on the sections.




Figure 23: X_0 after changes

Let X be the solid torus that we had at last in Figure 23. Then, we have

$$\langle \pi_j, \pi_j \rangle_{Z(S^1 \times S^1)} = \sum_{\xi_k} \alpha(\xi_k) Z_\Delta(X), \quad (4.4)$$

where $\alpha(\xi_k)$'s are positive coefficients determined by some products of statistical dimensions. It is obvious that each $Z_\Delta(X)$ is non-negative.

Since $S(p_j) = \sum_i S_{ij} p_i$, it follows from Lemma 4.5 that $Z_\Delta(D^2 \times S^1; S(p_j)) = S_{0j}$.

It is now easy to see that the value of the solid torus labeled by $S(p_j)$ has a similar expression of the summation when it is cut along the meridian. Namely, S_{0j} is written in the form $S_{0j} = \sum_{\xi_k} \beta(\xi_k) Z_\Delta(X)$, where $\beta(\xi_k)$'s are strictly positive coefficients determined by n_j and some products of statistical

dimensions. As we saw, not all of the values of $Z_\Delta(X)$ can be zero, so one of them must be strictly positive. Hence, $S_{0j} > 0$.

Condition (ii.a), (iii): Since both S and T -matrices are unitary, we check that S is symmetric and T is diagonal with respect to $\{p_0, \dots, p_r\}$.

First we prove that S is symmetric. From a topological observation of the $Z(S)$ -action on $Z(S^1 \times S^1)$, we have

$$\begin{aligned} S_{ij} &= \langle S(p_i), p_j \rangle_{Z(S^1 \times S^1)} = \overline{\langle S(p_i)^*, p_j^* \rangle_{Z(S^1 \times S^1)}} = \overline{\langle S(p_i)^*, p_j \rangle_{Z(S^1 \times S^1)}} \\ &= \overline{\langle S^*(p_i), p_j \rangle_{Z(S^1 \times S^1)}} = \overline{\langle p_i, S(p_j) \rangle_{Z(S^1 \times S^1)}} = \langle S(p_j), p_i \rangle_{Z(S^1 \times S^1)} \\ &= S_{ji}. \end{aligned}$$

Next, we prove that T is diagonal. From a topological observation of the $Z(T)$ -action on $Z(S^1 \times S^1)$, we have

$$\langle T(X) \cdot T^*(Y), Z \rangle_{Z(S^1 \times S^1)} = \langle X \cdot Y, Z \rangle_{Z(S^1 \times S^1)}$$

for any $X, Y, Z \in V(S^1 \times S^1)$, where $\cdot$ stands for the multiplication in the tube algebra. Hence, we have $T(X) \cdot T^*(Y) = X \cdot Y$ for any $X, Y \in V(S^1 \times S^1)$. This implies that T is diagonal.

Condition (ii.c.2): (Verlinde identity) From the definition of the fusion algebra associated with the Turaev-Viro-Ocneanu TQFT, N_{ij}^k is a structure constant of the fusion algebra.

From the unitarity of S , it is enough to prove

$$\sum_{i,j} N_{ij}^k \overline{S_{im}} \overline{S_{jn}} = \delta_{mn} \frac{\overline{S_{mk}}}{S_{m0}}. \quad (4.5)$$

The first term of the left-hand side of the equation (4.5) is written as $Z_\Delta(D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1); p_i, p_j; p_k)$. The second and the third terms of the left-hand side of the equation (4.5) can be written as $Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; p_i, S(p_m))$, $Z_\Delta(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; p_j, S(p_n))$, respectively. Use gluing and embed the second and third terms into the first term, then the left-hand side of the equation (4.5) is equal to $Z_\Delta(D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1); S(p_m), S(p_n); p_k)$. Then, cut $D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1)$ along the meridian.




Figure 24:

By using the gluing axiom of TQFT, we separate the rectangular part of this cut object by taking the summation of Z_Δ of the 3-manifold depicted in

Figure 24 over orthonormal basis η_i 's on the rectangle. Then, by the identity (4.3), we get the value 0 if $m \neq n$ because π_m and π_n are central orthogonal projections. Hence,

$$\begin{aligned}
& Z_{\Delta}(D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1); S(p_m), S(p_n); p_k) \\
= & \delta_{mn} \frac{\sum_m Z_{\Delta}(D^2 \times S^1 \setminus (\text{Int} D_1^2 \times S^1 \cup \text{Int} D_2^2 \times S^1); S(p_m), S(p_n); p_k) Z_{\Delta}(D^2 \times S^1; S(p_m))}{Z_{\Delta}(D^2 \times S^1; S(p_n))} \\
= & \delta_{mn} \frac{Z_{\Delta}(D^2 \times S^1 \setminus \text{Int} D_0^2 \times S^1; S(p_m), p_k)}{Z_{\Delta}(D^2 \times S^1; S(p_n))} \\
= & \delta_{mn} \frac{\overline{S_{mk}}}{S_{n0}}.
\end{aligned}$$

Condition (i): From the Verlinde identity and the unitarity of S , we have

$$N_{0j}^k = \sum_{l=0}^r \frac{S_{0l} S_{jl} \overline{S_{kl}}}{S_{0l}} = \sum_{l=0}^r S_{jl} \overline{S_{kl}} = \delta_{jk}.$$

So p_0 is the identity element in the fusion algebra.

Condition (ii.b): Since S^2 is $*$ -antiisomorphism, it is clear that $S^2(p_i)$'s are minimal projections again.

We have $S^2(p_0) = p_0$ because the definition of p_0 is invariant under the 180 degree rotation.

Condition (iv): We note that $Z(U)$ is nothing but the $*$ -operation of Tube Δ . So $Z(U)(p_i) = p_i^* = p_i$. The canonical map θ maps an orthonormal basis in $Z(-S^1 \times S^1)$ to the dual basis in $Z(S^1 \times S^1)^*$. So, $\theta(p_i) = \widehat{p}_i$, where $\{\widehat{p}_i\}_{i=0}^r$ is a dual basis of $\{p_i\}_{i=0}^r$ in $Z(S^1 \times S^1)^*$ such that $\widehat{p}_i(p_j) = \delta_{ij}$. Getting together, we have $\theta \circ Z(U)(p_i) = \theta(p_i) = \widehat{p}_i$. Q.E.D.

5 Applications

From the conclusion of Section 4, we know that there exists a Verlinde basis of $Z_{\Delta}(S^1 \times S^1)$ in the sense of Section 2 in a Turaev-Viro-Ocneanu TQFT. Hence, for a closed 3-manifold M , we have the following Dehn surgery formula:

$$Z_{\Delta}(M) = \sum_{i_1, \dots, i_m=0}^r S_{0i_1} \cdots S_{0i_m} J(L; i_1, \dots, i_m),$$

where we have assumed that the manifold M is obtained from S^3 by Dehn surgery along a framed link $L = L_1 \cup \cdots \cup L_m$.

The purpose in this section is to understand the formula of right-hand side of the above equation by introducing a notion of the *tube system* due to Ocneanu [14, 15], which gives a tensor category.

5.1 Tube systems

Let us start with the definition of a tube system. Let Δ be a finite system. A *tube system* $\mathcal{D}(\Delta)$ is a tensor category defined in the following way. First of all, the objects of $\mathcal{D}(\Delta)$ are the $\mathbb{C}$ -linear span of all minimal projections in the tube algebra $\text{Tube}\Delta$. For minimal projections p_i and p_j , the hom-set $\text{Hom}(p_i, p_j)$ is the set of vectors in the labeled surface depicted in Figure 25. For general objects $X, Y \in \mathcal{D}(\Delta)$, we define $\text{Hom}(X, Y)$ by extending $\text{Hom}(p_i, p_j)$ by linearity. Let $\sim$ be the Murray-von Neumann equivalence re-




Figure 25: $\text{Hom}(p_i, p_j)$

lation between projections. (Namely, two projections $p, q \in \text{Tube } \Delta$ are equivalent in the sense of Murray-von Neumann if there exists a partial isometry $v \in \text{Tube } \Delta$ such that $p = v^*v$ and $q = vv^*$.) If $p_i \sim p_j$, then $\text{Hom}(p_i, p_j) \cong \mathbb{C}$ and if $p_i \not\sim p_j$, then $\text{Hom}(p_i, p_j) = \{0\}$. Hence, minimal projections are simple objects in $\mathcal{D}(\Delta)$. We denote $\text{Hom}(p_i, p_i)$ by $\text{End}(p_i)$ for simplicity.

For simple objects p, q and r , the composition $x \cdot y$ of $x \in \text{Hom}(p, q)$ and $y \in \text{Hom}(q, r)$ is defined by the concatenation of x followed by y . See Figure 26. For general objects p, q and r , we define the composition of two morphisms as above by linearity. We also define x^* for x in $\text{Hom}(p, q)$ by inverting x inside out.

$$\begin{array}{c} \text{---} x \text{---} \\ p \quad q \end{array} \bullet \begin{array}{c} \text{---} y \text{---} \\ q \quad r \end{array} = \begin{array}{c} \text{---} x \text{---} y \text{---} \\ p \quad q \quad r \end{array}$$

Figure 26: The composition of $x \in \text{Hom}(q, r)$ and $y \in \text{Hom}(p, q)$

Let p_j be a simple object and q be an object in $\mathcal{D}(\Delta)$. We define an inner product $\langle x, y \rangle$ of $x, y \in \text{Hom}(q, p_i)$ by $\langle x, y \rangle = x \cdot y^*$, as a composition of morphisms. Then, with this inner product, $\text{Hom}(q, p_i)$ becomes a Hilbert space.

Next, we define the tensor product $p_i \otimes p_j$ of two simple objects p_i and p_j by the fusion product $p_i * p_j$ which was defined in Section 2. Then,

$\text{Hom}(p_i \otimes p_j, p_k)$ consists of vectors in the labeled surface of Figure 22. We define an inner product $\langle x, y \rangle$ of $x, y \in \text{Hom}(p_i \otimes p_j, p_k)$ by a composition of two morphisms $\langle x, y \rangle = x \cdot y^* \in \text{End}(p_k) \cong \mathbb{C}$. (Here, y^* is defined by inverting y (pants) inside out.) Note that we have Frobenius reciprocities for morphisms of $\text{Hom}(p_i \otimes p_j, p_k)$. For instance, we obtain $\text{Hom}(p_i \otimes p_j, p_k) \cong \text{Hom}(p_i, p_k \otimes p_{\bar{j}})$. See Figure 27. Frobenius reciprocities are given by the graphical operations in the category $\mathcal{D}(\Delta)$.




Figure 27: A Frobenius reciprocity map

We call the above defined semi-simple tensor category $\mathcal{D}(\Delta)$ a *tube system*.

To compute the quantum dimension of p_i , we make the composition of morphisms $b_i^* \circ b_i$, where b_i is the distinguished morphism from p_0 to $p_i \otimes p_{\bar{i}}$ obtained from Frobenius reciprocity of $\text{id}_{p_i} \in \text{End}(p_i)$. It must be a scalar multiple of p_0 since p_0 is a simple object. We put this value c . (See Figure 28.)




Figure 28: Computing a quantum dimension

To obtain the value c , we connect the upper p_0 and the lower p_0 , and embed it into S^3 . Fill out the outside of the tube, and take the Turaev-Viro-Oceanu invariant of it. Then, we get the $c = J(\bigcirc; i)/S_{00}$. This is the quantum dimension of p_i , denoted by $\dim p_i$.

We define the map c_{p_i, p_j} from $p_i \otimes p_j$ to $p_j \otimes p_i$ as in Figure 29. It is easy to see that this map c_{p_i, p_j} satisfies the axioms of a braiding, since the map c_{p_i, p_j} is defined in a topological way. So, we now know the tensor category $\mathcal{D}(\Delta)$ is braided.

We further define a map $\theta_{p_i} \in \text{End}(p_i)$ by Figure 30, where Z_Δ is evaluated at the 3-ball removed a twisted solid tube and the two tubes in the

$$c_{p_i, p_j} = \sum_{\xi_m, \eta_n, k} Z_{\Delta} \left(\text{Diagram 1} \right) \text{Diagram 2}$$

Figure 29: A map c_{p_i, p_j}

boudaries of it are both labeled by p_i . This map satisfies the following two

$$\theta_{p_i} = Z_{\Delta} \left(\text{Diagram 1} \right) \text{Diagram 2}$$

Figure 30: The twist map

equalities. (By the definition of θ_{p_i} , these equalities are proven by making pictures corresponding to the formulas in both sides and using some topological moves.)

$$\begin{aligned} \theta_{p_i \otimes p_j} &= c_{p_j, p_i} \circ c_{p_i, p_j} \circ (\theta_{p_i} \otimes \theta_{p_j}), \\ (\theta_{p_i} \otimes id_{p_i}) \circ b_i &= (id_{p_i} \otimes \theta_{p_i}) \circ b_i. \end{aligned}$$

Namely, θ_{p_i} defines a twist on $\mathcal{D}(\Delta)$ and this makes $\mathcal{D}(\Delta)$ a ribbon category.

Let us make the following compositions of morphisms in $\mathcal{D}(\Delta)$.

$$\begin{aligned} &b_0 \circ (id_{p_0} \otimes b_j) \circ (b_i \otimes id_{p_j} \otimes id_{b_j}) \circ (id_{p_i} \otimes c_{p_i, p_j} \otimes id_{p_j}) \circ (id_{p_i} \otimes c_{p_j, p_i} \otimes id_{p_j}) \\ &\circ (b_i^* \otimes id_{p_j} \otimes p_j) \circ (id_{p_0} \otimes b_j^*) \circ b_0^* \end{aligned}$$

Then, it is a scalar multiple of p_0 and makes a Hopf link H as a diagram. (See Figure 31.) Let us denote this scalar of the Hopf link H by s_{ij} . Embed this compositions of tubes into S^3 and fill out the outside of tubes. By Proposition 2.4 in Section 2, this provides

$$s_{ij} = \frac{J(H; i, j)}{S_{00}} = \frac{S_{ij}}{S_{00}},$$

where $S = (S_{ij})_{i, j=0}^r$ is the S -matrix with respect to our Verlinde basis $\{p_0, \dots, p_r\}$. Since the S -matrix is unitary, the matrix $(s_{ij})_{i, j=0}^r$ is invertible. It means that our category $\mathcal{D}(\Delta)$ is modular.

Remark 5.1. *The notions of the tube algebra and the tube system are also described by the language of a II_1 -subfactor $N \subset M$ with finite Jones index and finite depth, although our exposition here uses an infinite subfactor.*




Figure 31: The Hopf link

5.2 Dehn surgery formula as a Reshetikhin-Turaev invariant

Let $\mathcal{C}$ be a modular category and $\{V_i\}_{i=0}^r$ its simple objects. Put $\Delta = \Delta_+ = \sum_{i=0}^r t_i^{-1}(\dim V_i)^2$, $\Delta_- = \sum_{i=0}^r t_i(\dim V_i)^2$ and $D = (\sum_{i=0}^r (\dim V_i)^2)^{1/2}$. (Here, for Δ , we followed the notation in [21].) Let M be a closed 3-manifold obtained from S^3 by Dehn surgery along a framed link $L = L_1 \cup \dots \cup L_m$. Then, the Reshetikhin-Turaev invariant of M is given by the formula

$$\tau(M) = \Delta^{\sigma(L)} D^{-\sigma(L)-m-1} \sum_{\lambda \in \text{Col}(L)} \left(\prod_{n=1}^m \dim V_{\lambda(n)} \right) F(L, \lambda),$$

where $\sigma(L)$ is the signature of L and $F(L, \lambda)$ is the invariant of the colored framed link (L, λ) . (See [21] for the details. Also see [16].)

We note that in the above formula $\dim V_{\lambda(n)} = s_{0\lambda(n)}$. Hence the original Reshetikhin-Turaev formula can be rewritten with the s -matrix in the form

$$\tau(M) = \Delta^{\sigma(L)} D^{-\sigma(L)-m-1} \sum_{\lambda \in \text{Col}(L)} \left(\prod_{n=1}^m s_{0\lambda(n)} \right) F(L, \lambda).$$

Now, we start with the modular category $\mathcal{D}(\Delta)$ defined in Section 5.1 instead of a general modular category and make the Reshetikhin-Turaev formula. In our case, we already have S -matrix $(S_{ij})_{i,j=0}^r$ from the Turaev-Viro-Oceanu TQFT, which is expressed with respect to a Verlinde basis $\{p_0, \dots, p_r\}$ in $Z_\Delta(S^1 \times S^1)$, and since we know that $\dim p_i = S_{0i}/S_{00}$, by using $D = 1/S_{00}$, we can rewrite the Reshetikhin-Turaev formula in the form

$$\tau(M) = \Delta^{\sigma(L)} D^{-\sigma(L)} \sum_{\lambda \in \text{Col}(L)} \left(\prod_{n=1}^m S_{0\lambda(n)} \right) S_{00} F(L, \lambda).$$

We will prove that $D = \Delta$. First of all, from the equality $Z_\Delta(S^3) = Z_\Delta(L(1, 1)) = \sum_{i=0}^r t_i^{-1} S_{0i}^2$, we have $\frac{1}{S_{00}} = \lambda = \sum_{i=0}^r t_i^{-1} S_{0i}^2$ in our notation. From this, we get

$$\frac{1}{S_{00}} = \lambda = \sum_{i=0}^r t_i^{-1} \left(\frac{S_{0i}}{S_{00}} \right)^2 = \sum_{i=0}^r t_i^{-1} (\dim p_i)^2 = \Delta_+.$$

Taking the complex conjugation of the above formula, we get

$$\lambda = \sum_{i=0}^r t_i (\dim p_i)^2 = \Delta_-.$$

Thus $\lambda^2 = \Delta_+ \Delta_- = \sum_{i=0}^r (\dim p_i)^2$. Namely, we have $D = \Delta$. So, the Reshetikhin-Turaev invariant constructed from $\mathcal{D}(\Delta)$ is given by

$$\tau(M) = \sum_{\lambda \in \text{Col}(L)} \left(\prod_{n=1}^m S_{0\lambda(n)} \right) S_{00} F(L, \lambda).$$

Since F is uniquely determined by the category of the ribbon tangles [21, Part I, Chapter I], taking the normalization into consideration, we have $F(L, \lambda) = \frac{J(L, \lambda)}{S_{00}}$. Namely, in our case,

$$\tau(M) = \sum_{\lambda \in \text{Col}(L)} \left(\prod_{n=1}^m S_{0\lambda(n)} \right) J(L, \lambda),$$

which is nothing but our Dehn surgery formula Proposition 2.3. Let us summarize this argument as a theorem.

Theorem 5.2. *Let Δ be a finite system. For a closed oriented 3-manifold M , the Reshetikhin-Turaev invariant $\tau(M)$ constructed from a tube system $\mathcal{D}(\Delta)$ coincides with the Turaev-Viro-Ocneanu invariant $Z_\Delta(M)$.*

The following corollary has been proven by several authors [21], [17] etc in various settings.

Corollary 5.3. *Let Δ be a finite C^* -tensor category arising from a subfactor. If Δ is a modular category, then $\mathcal{D}(\Delta)$ is equivalent to $\Delta \otimes \Delta^{\text{op}}$ and this provides us with $\tau(M) \cdot \overline{\tau(M)} = Z_\Delta(M)$ for a closed oriented 3-manifold M , where τ is the Reshetikhin-Turaev invariant constructed from Δ .*

Proof. In [4], it is proved that $\mathcal{D}(\Delta)$ is equivalent to $\Delta \otimes \Delta^{\text{op}}$ when Δ is a modular category. (See Appendix too.) Hence, the rest is clear from Theorem 5.2. Q.E.D.

A Appendix

In this Appendix, we fix inaccuracies in [2] and [3]. In [2] and [3], the authors have analyzed the structure of M_∞ - M_∞ bimodules obtained from the asymptotic inclusion $M \vee M^{\text{op}} \subset M_\infty$ starting from the inclusion of AFD II_1 factors $N \subset M$ with finite Jones index and finite depth, following Ocneanu.

Let $\{p_0, \dots, p_r\}$ be a representative set of the equivalence classes of minimal projections of $\text{Tube}\Delta$ by the Murray-von Neumann equivalence relation.




Figure 32: The bimodule $X(p_i)$

We present the M_∞ - M_∞ bimodule $X(p_i)$ by Figure 32, where the circle at the middle is empty and the annulus around it is labeled with the minimal projection p_i . (See Chapter 12 in [2] for more explanation of this kind of pictures.) We can prove that each M_∞ - M_∞ bimodule $X(p_i)$ ($i = 0, \dots, r$) is irreducible in a similar way to the proof of Theorem 12.26 in [2].

Since $[_{M_\infty} X(p_i)_{M_\infty}]^{1/2} = \dim p_i$, we have the equalities $\sum_{i=0}^r [X(p_i)] = \sum_{i=0}^r (\dim p_i)^2 = \lambda^2$, where $\dim p_i$ is the quantum dimension of p_i as an object of $\mathcal{D}(\Delta)$. On the other hand, the global index of the asymptotic inclusion is given by λ^2 . This means that all the irreducible M_∞ - M_∞ bimodules obtained from the asymptotic inclusion are given by $\{X(p_i)\}_{i=0}^r$.

Remark A.1. In [2] and [3], irreducible bimodules are labeled by π_i 's, i.e., minimal central projections of $\text{Tube}\Delta$, instead of the minimal projections of it, which is incorrect.

We describe the fusion rule of irreducible M_∞ - M_∞ bimodules. The relative tensor product of two M_∞ - M_∞ bimodules $X(p_i) \otimes_{M_\infty} X(p_j)$ is decomposed into irreducible bimodules as in Figure 33.




Figure 33: Decomposition of a bimodule

Hence, the fusion rule is given by the fusion rule of the fusion algebra associated with the Turaev-Viro-Ocneanu TQFT.

It is easy to see that M_∞ - M_∞ bimodules arising from the asymptotic inclusion $M \vee M^{\text{op}} \subset M_\infty$ give rise to a modular category with the same braiding and twist as ones in a tube system. We denote this category of M_∞ - M_∞ bimodules by $\mathcal{M}_\infty$. To a simple object $X(p_i)$ in $\mathcal{M}_\infty$, we assign a simple object p_i in $\mathcal{D}(\Delta)$. From Figure 33, it is easy to see that we have a functor $F : \mathcal{M}_\infty \longrightarrow \mathcal{D}(\Delta)$, when we look at the morphisms in $\mathcal{M}_\infty$.

Let us now consider the opposite direction to the functor F . For a given morphism $x \in \text{Hom}(p_i \otimes p_j, p_k)$ of $\mathcal{D}(\Delta)$, we will construct a homomorphism in $\text{Hom}(X(p_i) \otimes_{M_\infty} X(p_j), X(p_k))$.




Figure 34: The bimodule $X(p_i) \otimes_{M_\infty} X(p_i)$

Let y_0 be an arbitrary element of $X(p_i) \otimes_{M_\infty} X(p_j)$. This y_0 is considered as an element in the bimodule in Figure 34. We denote it by y_1 . Then, we attach $x \in \text{Hom}(p_i \otimes p_j, p_k)$ to this y_1 by using two tubes p_i, p_j . See Figure 35. The central part of Figure 35 can be viewed as in Figure 36. Hence, we get a morphism in $\text{Hom}(X(p_i) \otimes_{M_\infty} X(p_j), X(p_k))$. This induces a functor $G : \mathcal{D}(\Delta) \longrightarrow \mathcal{M}_\infty$. It is not difficult to see that F and G are functors, which preserve the operations in modular categories, and inverse to each other.

Hence, the category of M_∞ - M_∞ bimodules obtained from the asymptotic inclusion $\mathcal{M}_\infty$ is equivalent to the tube system $\mathcal{D}(\Delta)$ constructed from $N \subset M$ as modular categories.




Figure 35:




Figure 36:

References

- [1] M. F. Atiyah, *Topological quantum field theories*, Publ. Math. I.H.E.S. 68 (1989) 175—186.
- [2] D. E. Evans, Y. Kawahigashi, *Quantum symmetries on operator algebras*, Oxford University Press, 1998.
- [3] D. E. Evans, Y. Kawahigashi, *On Ocneanu’s theory of asymptotic inclusions for subfactors, topological quantum field theories and quantum doubles*, Internat. J. Math. 6 (1995) 205–228.
- [4] D. E. Evans, Y. Kawahigashi, *Orbifold subfactors from Hecke algebras II*, Commun. Math. Phys. 196 (1998) 331–361.
- [5] L. Feher, A. Stipsicz and J. Szenthe ed., *Topological field theories and geometry of loop spaces*, World Scientific, 1992.
- [6] M. Izumi, *Applications of fusion rules to classification of subfactors*, Publ. RIMS, 27, (1991), 953–994.
- [7] M. Izumi, *The structures of sectors associated with the Longo-Rehren inclusions I. General theory*, Commun. Math. Phys. 213 (2000) 127—179.
- [8] M. Izumi, *The structures of sectors associated with the Longo-Rehren inclusions II. Examples*, Rev. in Math. Phys. 13 (2001) 603—674.
- [9] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, Bull. of the AMS, 12 (1985), 103–112.
- [10] T. Kerler, *Genealogy of non-perturbative quantum-invariants of 3-manifold: the surgical family*, in “Geometry and Physics” (Aarhus 1995), 503–547. Lect. Notes in Pure and Appl. Math., 184, Dekker, New York, 1997.
- [11] T. Kohno, *Topological invariants for 3-manifolds using representations of mapping class groups II: Estimating tunnel number of knots*, Contemp. Math. 175 (1994) 193–217.
- [12] R. Longo and J. E. Roberts, *A theory of dimension, K-theory* 11 (1997), 103–159.
- [13] M. Müger, *From subfactors to categories and topology II. The quantum double of subfactors and categories*, preprint, math.CT/0111205, 2001.
- [14] A. Ocneanu, *Chirality for operator algebras*, 39—63, in “Subfactors”, ed. by H. Araki, et al., World Scientific, 1994.

- [15] A. Ocneanu, *Operator Algebras, Topology and Subgroups of Quantum Symmetry—Construction of Subgroups of Quantum Groups—*, 235—263, written by Satoshi Goto(Sophia University) and Nobuya Sato(Osaka Prefecture University), in “Advanced Studies in Pure and Mathematics” 31, Taniguchi Conference in Mathematics Nara '98, Mathematical Society of Japan, 2000.
- [16] N. Reshetikhin and V. G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. 103 (1991), p. 547–597.
- [17] J. Roberts, *Skein theory and Turaev-Viro invariants*, Topology 34 (1995), 771–787.
- [18] N. Sato and M. Wakui, *(2+1)-dimensional topological quantum field theory with Verlinde basis and Turaev-Viro-Ocneanu invariants of 3-manifolds*, preprint 2000.
- [19] K. Suzuki and M. Wakui, *On the Turaev-Viro-Ocneanu invariant of 3-manifolds derived from the E_6 -subfactor*, Kyushu J. Math. 56 (2002) 59–81.
- [20] T. Takata, *Invariants of 3-manifolds associated with quantum groups and Verlinde’s formula*, Publ. RIMS Kyoto Univ. 28 (1992) 139–167.
- [21] V. G. Turaev, *Quantum invariants of knots and 3-manifolds*, Walter de Gruyter 1994.
- [22] V. G. Turaev and O. Ya Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, Topology 31 (1992) 865—902.
- [23] E. Verlinde, *Fusion rules and modular transformations in 2D conformal field theory*, Nucl. Phys. B300 (1988) 360–376.
- [24] M. Wakui, *Fusion algebras for orbifold models (a survey)*, in “Topology, geometry and field theory” edited by K. Fukaya, M. Furuta, T. Kohno and D. Kotschick, World Scientific, 1994, 225–235.
- [25] E. Witten, *Quantum field theory and Jones polynomial*, Commun. Math. Phys. 121 (1989) 351–399.

